

## SNAPSHOT OVERVIEW OF NEW QM / QI REPORTS

Measure Description	Specifications
<b>Accidents</b>	
<b>1.1</b> Incidence of new fractures	New hip fracture or other fracture checked on target assessment and not checked on prior assessment
<b>1.2</b> Prevalence of falls	Residents who had falls within the past 30 days
<b>Behavior/Emotional Patterns</b>	
<b>2.1</b> Residents who have become more depressed or anxious	Mood scale scores on target assessment are > than scores on prior assessment
<b>2.2</b> Prevalence of behavior symptoms affecting others	Residents with behavior symptoms affecting others (verbally abusive, physically abusive, or socially inappropriate/disruptive) on target assessment
<b>2.2</b> High-risk	Those who are cognitively impaired, have psychotic disorders, or are manic-depressive
<b>2.2</b> Low-risk	All others
<b>2.3</b> Prevalence of symptoms of depression without antidepressant therapy	Residents with symptoms of depression and no antidepressant therapy on target assessment
<b>Clinical Management</b>	
<b>3.1</b> Use of 9 or more different medications	Residents who received 9 or more different medications on target assessment
<b>Cognitive Patterns</b>	
<b>4.1</b> Incidence of cognitive impairment	Residents who were cognitively impaired on target assessment (short-term memory problems and any impairment in daily decision making)
<b>Elimination/Incontinence</b>	
<b>5.1</b> Low-risk residents who lost control of their bowels or bladder	Residents who were frequently or fully incontinent on target assessment
<b>5.2</b> Residents who have/had a catheter inserted and left in their bladder	Residents with indwelling catheters on target assessment
<b>5.3</b> Prevalence of occasional or frequent bladder or bowel incontinence without a toileting plan	Residents with no scheduled toileting plan and no bladder-retraining program on target assessment
<b>5.4</b> Prevalence of fecal impaction	Residents with fecal impaction on target assessment
<b>Infection Control</b>	
<b>6.1</b> Residents with a urinary tract infection (UTI)	Residents with a UTI on target assessment
<b>Nutrition/Eating</b>	
<b>7.1</b> Residents who lose too much weight	Residents with 5% weight loss in last 30 days, or 10% weight loss in last 6 months
<b>7.2</b> Prevalence of tube-feeding	Residents with a tube-feeding on target assessment
<b>7.3</b> Prevalence of dehydration	Residents with dehydration on target assessment (output exceeds intake, or ICD-9 code 276.5)

<b>Pain Management</b>	
<b>8.1</b> Residents who have moderate to severe pain	Residents with moderate pain at least daily, or horrible/excruciating pain at any frequency on target assessment
<b>Physical Functioning</b>	
<b>9.1</b> Residents whose need for help with daily activities has increased	Late-loss ADLs have worsened (i.e., bed mobility, transfer, eating, and toileting)
<b>9.2</b> Residents who spend most of their time in a bed or in a chair	Residents who a bedfast on target assessment
<b>9.3</b> Residents whose ability to move around in their room got worse	Residents whose value for locomotion is greater on target assessment than on prior assessment
<b>9.4</b> Incidence of decline in ROM	Residents with increases in functional limitation in ROM between prior and target assessments
<b>Psychotropic Drug Use</b>	
<b>10.1</b> Prevalence of antipsychotic use, in the absence of psychotic or related conditions	Residents receiving antipsychotics with no psychotic or related disorder or diagnosis
<b>10.1</b> High-risk	Includes cognitive impairment and behavior problems
<b>10.1</b> Low-risk	Includes all others
<b>10.2</b> Prevalence of antianxiety/hypnotic use	Residents who received antianxiety or hypnotic medications on target assessment.
<b>10.3</b> Prevalence of hypnotic use more than 2 times in last week	Residents who received hypnotics > 2 times in last week on target assessment
<b>Quality of Life</b>	
<b>11.1</b> Residents who were physically restrained	Residents who were physically restrained daily on target assessment (does not include side rails)
<b>11.2</b> Prevalence of little or no activity	Residents with little or no activity on target assessment
<b>Skin Care</b>	
<b>12.1</b> High-risk residents with pressure ulcers	Residents with Stage 1–4 pressure ulcers and have impaired bed mobility or transfers, are comatose, or have malnutrition on target assessment
<b>12.2</b> Low-risk residents with pressure ulcers	All others
<b>Post-Acute Care (PAC) Measures</b>	
<b>13.1</b> Short-stay residents with delirium	Short-stay residents with at least 1 symptom of delirium on 14-day PPS assessment
<b>13.2</b> Short-stay residents who had moderate to severe pain	Short-stay residents with moderate pain at least daily or horrible/excruciating pain at any frequency on 14-day PPS assessment
<b>13.3</b> Short-stay residents with pressure ulcers	Short-stay residents who had no pressure sores on PPS 5-day assessment and had at least a stage 1 pressure sore on the 14-day PPS assessment, OR Who had a pressure sore on the PPS 5-day assessment, that worsened or failed to improve on the 14-day PPS assessment

Source: Sylvia Balistreri, RN, BSN, Program Manager, Office of Long Term Care Licensing, ADHS