

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Accidents</i>			
1.1 Incidence of new fractures	1		
1.2 Prevalence of falls	2		
<i>Behavior/Emotional Patterns</i>			
2.1 Residents who have become more depressed or anxious		CMOD03	Replaces QI 4 (prevalence of symptoms of depression). Note: QI 4 was a prevalence measure, while the new measure is an incidence measure and may have quite different values.
2.3 Prevalence of symptoms of depression without antidepressant therapy	5		

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Behavior/Emotional Patterns</i>			
2.2 Prevalence of behavior symptoms affecting others	3		
2.2-HI High risk	3-HI		
2.2-LO Low risk	3-LO		
<i>Clinical Management</i>			
3.1 Use of 9 or more different medications	6		
<i>Cognitive Patterns</i>			
4.1 Incidence of cognitive impairment	7		

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Elimination/Incontinence</i>			
5.1 Low-risk residents who lost control of their bowels or bladder		CCNT06	Replaces QI 8-LO (prevalence of bladder or bowel incontinence: low risk)
5.3 Prevalence of occasional or frequent bladder or bowel incontinence without a toileting plan	9		
5.2 Residents who have/had a catheter inserted and left in their bladder		CCAT02	Replaces QI 10 (prevalence of indwelling catheter)
5.4 Prevalence of fecal impaction	11		
<i>Infection Control</i>			
6.1 Residents with a urinary tract infection		CCNT04	Replaces QI 12 (prevalence of urinary tract infection)

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Nutrition/Eating</i>			
7.1 Residents who lose too much weight		CWLS	Replaces QI 13 (prevalence of weight loss) and differs from QI 13 as an exclusion for residents with hospice care has been added.
7.2 Prevalence of tube feeding	14		
7.3 Prevalence of dehydration	15		
<i>Pain Management</i>			
8.1 Residents who have moderate to severe pain		CPAIOX	

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Physical Functioning</i>			
9.1 Residents whose need for help with daily activities has increased		CADL01	Replaces QI 17 (incidence of decline in late loss ADLs)
9.2 Residents who spend most of their time in bed or in a chair		CBFT01	Replaces QI 16 (prevalence of bedfast residents).
9.3 Residents whose ability to move in and around their room got worse		CMOB01	
9.4 Incidence of decline in ROM	18		
<i>Psychotropic Drug Use</i>			
10.1 Prevalence of antipsychotic use, in the absence of psychotic or related conditions	19		
10.1-HI High risk	19-HI		

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Psychotropic Drug Use</i>			
10.1-LO Low risk	19-LO		
10.2 Prevalence of antianxiety/hypnotic use	20		
10.3 Prevalence of hypnotic use more than two times in last week	21		
<i>Quality of Life</i>			
11.1 Residents who were physically restrained		CRES01	Replaces QI 22 (prevalence of daily physical restraints)
11.2 Prevalence of little or no activity	23		

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.

Measure Comparison - New Versus Old Measures

QI/QM Measures on New Reports	QI Measure ¹	QM Measure ²	Comments ³
<i>Chronic Care Measures</i>			
<i>Skin Care</i>			
12.1 High-risk residents with pressure ulcers		CPRU02	Replaces QI 24-HI (prevalence of stage 1-4 pressure ulcers: high risk)
12.2 Low-risk residents with pressure ulcers		CPRU03	Replaces QI 24-LO (prevalence of stage 1-4 pressure ulcers: low risk)
<i>Post-Acute Care (PAC) Measures</i>			
13.1 Short-stay residents with delirium		PAC-DEL0X	
13.2 Short-stay residents who had moderate to severe pain		PAC-PAI0X	
13.3 Short-stay residents with pressure ulcers		PAC-PRU0X	

¹QI numbers correspond to the numbering on the old QI reports.

²QM abbreviations correspond to their designations in the QM user's manual.

³ QMs have replaced QIs when their definitions are similar (although they may not be identical). QIs with no equivalent QM have been retained. Three QIs have been dropped without replacement: QI 8 [Overall] Prevalence of bladder or bowel incontinence, QI 8-HI High risk prevalence of bladder or bowel incontinence, and QI-24 [Overall] Prevalence of stage 1-4 pressure ulcers.